


**ADVANCE SOCIAL SCIENCE ARCHIVE JOURNAL**

 Available Online: <https://assajournal.com>

Vol. 05 No. 01. Jan-March 2026. Page#.1331-1337

 Print ISSN: [3006-2497](https://doi.org/10.5281/zenodo.18709199) Online ISSN: [3006-2500](https://doi.org/10.5281/zenodo.18709199)

 Platform & Workflow by: [Open Journal Systems](https://doi.org/10.5281/zenodo.18709199)
<https://doi.org/10.5281/zenodo.18709199>


## Influence of Body Parts Measurements with Low and High Risk of Obesity Using Male Human Body Fat Percentage

**Mir Ghulam Hyder Talpur**

Department of Mathematics and Statistics, Institute of Business Management, Karachi

[hyder.talpur@iobm.edu.pk](mailto:hyder.talpur@iobm.edu.pk)

0301-3409885

**Muhammad Shahbaz Khan**

Department of Mathematics and Statistics, Institute of Business Management, Karachi

[shahbazk@kiet.edu.pk](mailto:shahbazk@kiet.edu.pk)

03222515941

### Abstract

*There are many indicators of the body that inform about the health of a person. In these indicators, Body-Fat-Percentage (BFP) is an important factor that classifies the low and high risk of obesity. Obesity is one of the silent killer illnesses that make people lazy and unable to perform their daily routine work smoothly. BFP factor distinguishes the status of normal and obese persons. In this study, we identified that all body measurements including density and age have significant differences between Low and High risk of obesity. Human body measurement factors are density (den), age, weight (w), height (h), neck (n), chest (c), abdomen (abd), hip, thigh (t), knee (K), ankle (A), biceps (b), forearm (f), and wrist (w) body measures. Screening of data was performed by removing outliers using Box-Whisker plots from all factors. Box-Whisker plot constructed through R software using package ggplot2. Comparison of body parts measurements based on Low-Risk and High-Risk using two-sample t-tests done through SPSS software.*

**Keywords:** *Body fat percentage, density, two-sample t-test, heatmap, obesity*

### Introduction

In medical science, obesity term is considered an increasingly chronic disease that leads to life-threatening problems such as heart attack or diabetes problems. A peculiar or high body fat buildup impairs health issues (Ellulu, et.al, 2017).

Obesity is the unhealthy status of a person whose body is at high risk of different diseases. It upsets the normal function of the heart, liver, kidneys, joints, and reproductive system (Fatima, et.al, 2023). It starts with a range of noncontagious diseases (NCDs), for example, Type-2 diabetes, heart and blood vessel disorders, and hypersensitivity. and increased chances of stroke, different types of cancer, as well as psychological disorders issues (Lin. C. L. et.al, 2021).

A person with obesity is three times more chance to be hospitalized for COVID-19 [WHO]. The global incidence of obesity has increased from 7% in 1980 to 13% in 2015, an increase of almost 85%, reaching epidemic levels (Petraakis, D., et.al, 2020). In medical science, BFP is a measurement that defines the status of a person as either obese or normal. BFP calculates how much of a person's body consists of fat (Chiplonkar et al., 2017). It helps doctors to recognize healthy and obese people more effectively.

Physical fitness and aerophilic exercises have significantly correlated with BFP (Shoebuddin & Daimi, 2019).

Non-alcoholic fatty liver disease (NAFLD) is rising internationally as the number of obesity cases increases. Among those cases who have NAFLD, a strong relationship between Low Muscle Mass (LMM) (e.g. sarcopenia) and high risk of NAFLD and fibrosis were observed with also some studies of child obesity (Y.M. Choi et.al, 2023). Obesity is the growth of peculiar or unnecessary body fat (BF) that may cause an unhealthy status. Controlling obesity decreases the chance of cardiovascular and stroke diseases (Akil, L., 2011).

A high correlation is observed between Type 2 diabetes mellitus (T2DM) with obesity, which can also be determined through Body Mass Index (BMI). It is also a fact that BMI cannot differentiate muscle mass and fat mass. Research shows that BMI is not a fat distribution in the body, which is a basic limitation. Studies showed that T2DM has a higher correlation between BFP and skeletal muscle than BMI (Bays, H. E., 2007). This study also mitigates the risks of global obese people (Safaei et al., 2021).

This study described the high-risk obesity in Karachi due to overweight and obesity dominance. A high rate of obesity is found in males as compared to females in Karachi. Some factors identified in this study such an unmoving lifestyle, unhealthy diet, and lack of physical workout. Preemptive actions regarding overweight and obesity may be suggested for people (Ibrahim et al., 2021).

A researcher found differences in body composition between playing standard and age in males who played rugby games. Spotlight training and nutritional schemes that minimize fat mass growth (Geeson-Brown et al., 2020). In America, a team distinguished adults according to cardiovascular fitness (CVf), BFP, and BMI in the National Health and Nutrition Examination Survey (NHANES) data (Bunn et al., 2019). Short-term and long-term diseases like hypertension, some types of cancer, cardiovascular, Alzheimer's, asthma, osteoarthritis, metabolic syndrome, musculoskeletal disorders, gallbladder disease, obstructive sleep apnea, hypercholesterolemia, metabolic syndrome, and type 2 diabetes caused by obesity (Mohajan & Mohajan, 2023)

Thant Zin and his team members have determined the mean BMI and average BFP with body fat assessments for obesity awareness among people. They suggested that including BMI, BFP other factors will be examined to monitor obesity such as age, sex, race, nutrition, and body parts measurements which change over time (Zin et al., 2014).

According to Sema Can examine the impacts of physical activity which control BMI, and body fat. In this research, shared workouts in both genders presented a positive impact on BMI and BFP (Can et al., 2019). Kathryn E. Bradbury and her team identified that those people who are continuously involved in physical activities have a low BFP after observing BMI importance (Bradbury et al., 2017).

## Materials and Methods

In this research study, the body fat percentage variable was converted into two groups. These two groups are designed to define the high and the low risk of obesity. A body fat percentage when more than 25 percent then it is considered to be a person at high risk of obesity. Similarly, less than 25 percent of the Body fat percentage is considered to be a person who is at a low risk of obesity. After making two groups, we now compare low-risk and high-risk obese person body measurements using two sample "t" tests. Data screening is necessary for removing the outliers therefore R package "ggplot2" (Wickham H., (2016) was used to construct the Box-Whisker plot. Two sample t-tests will be performed through SPSS (IBM Corp. 2011) to compare the body parts measurements among low and high risk of obesity.

### A. Data set:

Secondary data is used to analyze the low-risk and high-risk obesity (Body Fat Percentage Dataset, 2023). This sample data consists of 252 men's information about body parts measurements including density, BFP, and age.

**B. Definitions:**

Some important definitions related to the study are as follows:

- (i) Obesity: Excess food and non-physical activity develop a chronic disease called Obesity. A series of extra food consumption forms excess calories which lead to cause of obesity (Mohajan & Mohajan, 2023).
- (ii) Body Fat Percentage: The quantity of fat mass observed on the body conveyed as the total mass of fat (TMoF) divided by total body mass (TBM), and multiplied by 100, i.e.,

$$BFP = \frac{\text{Total mass of fat}}{\text{Total body mas}} \times 100$$

Body fat percentage calculators can be used to measure the individual person's body fat percentage (Body Fat Calculator, 2023).

- (iii) Density: Density is also known as Hydrostatic weighing. According to it, the body's density is calculated by getting the change in body weight in air and underwater based on Archimedes Rule.
- (iv) Body Measurements: Body measurements such as neck, chest, abdomen, hip, and thigh. Knee and others are very important to compare between low-risk and high-risk obese persons.

Table 1 shows the body measurements with their units.

Table 1: *Body measurements with their units:*

Measurements	Units	Measurements	Units
Weight	lbs.	Thigh circumference	cm
Height	cm	Knee circumference	cm
Neck circumference	cm	Ankle circumference	cm
Chest circumference	cm	Biceps (extended) circumference	cm
Abdomen 2 circumference	cm	Forearm circumference	cm
Hip circumference	cm	Wrist circumference	cm

**Results & Discussion**


Statistical results help us to analyze the low and high risk of obesity in men. First, remove the outlier's values from the data. Out of 252, 232 values are left for analysis. Box and Whisker plots were used to find outliers from each factor. Secondly, determine basic statistics for checking the basic assumptions about the normality of the variables to apply the statistical method "Two sample t-test" presented in Table 2. Thirdly, a "two sample t-test" was applied to determine the difference between the two groups' low and high risk of obesity persons using different factors. Results are presented in Table 3. The following flow diagram shows the steps:

- (i) Applied Box-Whisker plot to remove outlier's values from the sample data.

- (ii) Compute the basic statistics to check the normality assumption for the two-sample t-test method.
- (iii) Applied two-sample t-test, over every factor with respect to groups with low and high risk of obesity.

Step 1: The Box-Whisker Plot after removing the extreme outliers from the data presented in Fig.1.

Figure 1: Box-Whisker Plot



Step 2: Compute the basic statistics of all variables to check the normality assumption for the two-sample t-test method presented in Table 2.

Table 2: Basic statistics for all variables:

Variables	Statistics							
	Average	Variance	Std. Deviation	Minimum	Maximum	Interquartile Range	Skewness	Kurtosis
Density	1.0569	0.0003	0.0179	1.0101	1.1089	0.0278	0.1633	-0.5173
BodyFat %	18.5459	61.5714	7.8467	0.0000	40.1000	12.3000	-0.0436	-0.6557
Age	44.8498	159.0937	12.6132	22.0000	81.0000	19.0000	0.2421	-0.4566
Weight	176.3135	605.8511	24.6140	118.5000	241.2500	35.8750	0.2484	-0.3811
Height	178.55	42.90	6.54	162.56	197.48	34.92	0.1200	-0.4010
Neck	37.8635	4.4880	2.1185	32.8000	43.9000	2.8500	0.1068	-0.3780
Chest	100.1554	56.0127	7.4842	79.3000	119.9000	10.6000	0.3764	0.0862
Abdomen	91.5790	85.7634	9.2609	69.4000	118.0000	14.2500	0.1781	-0.3696
Hip	99.2322	33.6927	5.8045	85.0000	114.3000	7.2000	0.2161	-0.0134
Thigh	59.0202	20.2887	4.5043	47.2000	71.2000	5.8500	0.1253	-0.1602
Knee	38.4361	4.6770	2.1626	33.4000	44.2000	2.8500	0.1592	-0.2051
Ankle	22.9296	1.6283	1.2760	19.1000	27.0000	1.8000	0.2937	0.0175
Biceps	32.1176	7.8794	2.8070	24.8000	39.1000	3.8500	0.1278	-0.3745
Forearm	28.6519	3.2018	1.7894	24.6000	33.7000	2.7000	0.1037	-0.4970
Wrist	18.1794	0.7041	0.8391	16.3000	20.4000	1.2000	0.0406	-0.4312

The summary of statistics shows that all variables follow the basic assumption of normality because skeweness lies within the -3 to +3 range. Now we can apply two sample t-tests for the comparison of variables between low and high risk of obese people.

Step 3: Two sample "t" test results are presented in Table 3:

**Table 3: T-test for Equality of Means**

Variables	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference		Results based on P value
Density	15.141	231	0	0.03	0.002	0.026	0.034	Important
BodyFat	-15.684	231	0	-13.326	0.85	-15	-11.652	Important
Age	-4.221	231	0	-7.981	1.891	-11.707	-4.256	Important
Weight	-6.528	231	0	-22.971	3.519	-29.905	-16.038	Important
Height	1.417	231	0.158	1.43	1.01	-0.5620	3.435	Not Important
Neck	-5.522	231	0	-1.711	0.31	-2.321	-1.1	Important
Chest	-8.479	231	0	-8.623	1.017	-10.626	-6.619	Important
Abdomen	-11.17	231	0	-12.969	1.161	-15.257	-10.682	Important
Hip	-6.685	231	0	-5.526	0.827	-7.155	-3.898	Important
Thigh	-5.284	231	0	-3.498	0.662	-4.802	-2.194	Important
Knee	-5.13	231	0	-1.636	0.319	-2.264	-1.007	Important
Ankle	-2.536	231	0.012	-0.497	0.196	-0.882	-0.111	Important
Forearm	-3.745	231	0	-1.012	0.27	-1.545	-0.48	Important
Wrist	-2.8	231	0.006	-0.359	0.128	-0.612	-0.106	Important

The two-sample t-test results show that all the factors are significantly playing role between low-risk obesity and high-risk obesity in men through two information P-value and Confidence interval of the differences except height. Height is the only factor which shows not important for the distinguish of low and high risk of obesity.

### Conclusions and Limitation

In this study, we compared different factors based on Low and High risk of obesity in male persons. Two sample t-tests methods compared the average of different factors and found significant differences between Low and High Risk of obesity personals. Researchers claimed in different studies found significant that physical workout is very important for maintaining a healthy life. A very important limitation of this study is only one gender was used for this analysis which is male. Future work would be possible with female gender and children also. Finally, body measurements such as hip, abdomen, weight, knee, and others played a significant role in low and high risk of obese male persons.

### Conflict of interest

There is no conflict of interest.

### Ethical information

This study is focused on the people who are involved and not involved in physical workouts for the betterment of their lives.

### Funding (if any)

There is no funding.

### Author contribution

Author contributions are as follows:

Author 1 conceived the idea and directions.

Author 2, a write-up of the whole paper.

Author 3, R program for graph.

Author4, data analysis.

### References:

- Akil, L., & Ahmad, H. A. (2011). Relationships between obesity and cardiovascular diseases in four southern states and Colorado. *Journal of health care for the poor and underserved*, 22(4 Suppl), 61–72. <https://doi.org/10.1353/hpu.2011.0166>
- Bays, H. E., Chapman, R. H., Grandy, S., & SHIELD Investigators' Group (2007). The relationship of body mass index to diabetes mellitus, hypertension and dyslipidaemia: comparison of data from two national surveys. *International journal of clinical practice*, 61(5), 737–747. <https://doi.org/10.1111/j.1742-1241.2007.01336.x>
- Body Fat Calculator*. (n.d.). Calculator.net. Retrieved September 6, 2023, from <https://www.calculator.net/body-fat-calculator.html>
- Body Fat Percentage*. (n.d.). Kaggle. Retrieved September 6, 2023, from <https://www.kaggle.com/code/mustafacihadgoktepe/body-fat-percentage>
- Bradbury, K. E. et al., (2017). Association between Physical Activity and Body Fat Percentage, with Adjustment for BMI: A Large CrossSectional Analysis of UK Biobank. *BMJ Open*, 7, e011843.
- Bunn, J. et al., (2019). Fitness and Fatness: Body Mass Index versus Percent Body Fat. *Journal of Clinical Exercise Physiology*, 8(4), 131-137.
- Can, S., Demirkan, E., & Erca, S., (2019). The Effects of Exercise Preferences on Body Fat and Body Mass Index by Self-Report. *Universal Journal of Educational Research*, 7(1), 293-297.
- Ellulu, M. S., Patimah, I., Khaza'ai, H., Rahmat, A., & Abed, Y. (2017). Obesity and inflammation: the linking mechanism and the complications. *Archives of medical science : AMS*, 13(4), 851–863. <https://doi.org/10.5114/aoms.2016.58928>
- Fatma, Ibrahim. Abdel-latif. Megahed., 2023. "Unpacking the Factors of Overweight/Obesity in Type 2 Diabetes: What You Need to Know: Review Article." *Sumerianz Journal of Medical and Healthcare*, vol. 2, pp. 8-12.
- Geeson-Brown, T., Jones, B., Till, K., Chantler, S., & Deighton, K., (2020). Body Composition Differences by Age and Playing Standard in Male Rugby Union and Rugby League: A Systematic Review and Meta-Analysis. *Journal of Sports Sciences*, <https://doi.org/10.1080/02640414.2020.1775990>.
- IBM Corp. Released 2011. *IBM SPSS Statistics for Windows, Version 20.0*. Armonk, NY: IBM Corp.
- Ibrahim, S., & Akram, Z. et al., (2021). Overweight and Obesity Prevalence and Predictors in People Living in Karachi. *Journal of Pharmaceutical Research International*, 33(31), 194-202
- Khadilkar, A., Chiplonkar, S., Kajale, N. et al. (2018). Impact of dietary nutrient intake and physical activity on body composition and growth in Indian children. *Pediatr Res* 83, 843–850 <https://doi.org/10.1038/pr.2017.322>
- Lin, C. L., Yu, N. C., Wu, H. C., Lee, Y. Y., Lin, W. C., Chiu, I. Y., Chien, W. C., & Liu, Y. C. (2021). Association of Body Composition with Type 2 Diabetes: A Retrospective Chart Review Study. *International journal of environmental research and public health*, 18(9), 4421. <https://doi.org/10.3390/ijerph18094421>
- Mohajan, Devajit & Mohajan, Haradhan, 2023. "Obesity and Its Related Diseases: A New Escalating Alarming in Global Health," MPRA Paper 116797, University Library of Munich, Germany, revised 05 Feb 2023.

- Mohajan, Devajit & Mohajan, Haradhan, 2023. "Obesity and Its Related Diseases: A New Escalating Alarming in Global Health," MPRA Paper 116797, University Library of Munich, Germany, revised 05 Feb 2023.
- Petrakis, D., Margină, D., Tsarouhas, K., Tekos, F., Stan, M., Nikitovic, D., Kouretas, D., Spandidos, D. A., & Tsatsakis, A. (2020). Obesity - a risk factor for increased COVID-19 prevalence, severity and lethality (Review). *Molecular medicine reports*, 22(1), 9–19. <https://doi.org/10.3892/mmr.2020.11127>
- Safaei, M., Sundararajan, E. A., Driss, M., Boulila, W., & Shapi'i, A. (2021). A systematic literature review on obesity: Understanding the causes & consequences of obesity and reviewing various machine learning approaches used to predict obesity. *Computers in biology and medicine*, 136, 104754. <https://doi.org/10.1016/j.compbiomed.2021.104754>
- Shoebuddin, M. & Daimi, . S. B. (2019) Correlation of percentage body fat with physical efficiency index and maximal oxygen uptake. *National Journal of Physiology, Pharmacy and Pharmacology*, 9 (7), 586-589. <https://doi:10.5455/njppp.2019.9.0312004042019>
- Wickham H (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. ISBN 978-3-319-24277-4, <https://ggplot2.tidyverse.org>.
- Y.M. Choi, J.M. Kim, H.R. Yang, (2023) Significance Of Fat-Free Mass Measurement Beyond Fat Mass In Obese Children With Non-Alcoholic Fatty Liver Disease. *Clinical Nutrition ESPEN*, Volume 54, 667, <https://doi.org/10.1016/j.clnesp.2022.09.612>.
- Zin, T. et al., (2014). Body Fat Percentage, BMI and Skinfold Thickness among Medical Students in Sabah, Malaysia. *South East Asia Journal of Public Health*, 4(1), 35-40.